

This is an electronic reprint of the original article.

This reprint *may differ* from the original in pagination and typographic detail.

Author(s): Lena Gustafsson, Mats Hannerz, Matti Koivula, Ekaterina Shorohova, Ilkka Vanha-Majamaa & Jan Weslien

Title: Research on retention forestry in Northern Europe

Year: 2020

Version: Published version

Copyright: The Author(s) 2020

Rights: CC BY 4.0

Rights url: <http://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Gustafsson, L., Hannerz, M., Koivula, M. et al. Research on retention forestry in Northern Europe. *Ecol Process* 9, 3 (2020). <https://doi.org/10.1186/s13717-019-0208-2>

All material supplied via *Jukuri* is protected by copyright and other intellectual property rights. Duplication or sale, in electronic or print form, of any part of the repository collections is prohibited. Making electronic or print copies of the material is permitted only for your own personal use or for educational purposes. For other purposes, this article may be used in accordance with the publisher's terms. There may be differences between this version and the publisher's version. You are advised to cite the publisher's version.

REVIEW

Open Access

Research on retention forestry in Northern Europe



Lena Gustafsson^{1*} , Mats Hannerz², Matti Koivula³, Ekaterina Shorohova^{4,5,6}, Ilkka Vanha-Majamaa⁵ and Jan Weslien⁷

Abstract

Retention approaches in forest management are today common in several North European countries, integrated into the clearcutting practice as a way to promote biodiversity and maintain ecosystem functions. Individual green trees and retention patches (tree groups) are retained at final harvest, and deadwood is left at site or created. Here, we review research on retention in Sweden, Finland, Norway, the Baltic States, and NW Russia, with special focus on biodiversity. Following the first publication in 1994, about 180 peer-reviewed articles have been published. We present results from a systematic search of the retention literature, separated into the following topics: buffer zones, retention patches, high stumps, other types of deadwood, European aspen *Populus tremula*, and cost-efficiency. Russian literature is synthesized separately since studies from this region have so far almost exclusively been published in the Russian language. Furthermore, we describe six ongoing large-scale, replicated experiments with varying retention levels, five in Finland and one in Sweden, and summarize their main results. Among main conclusions for practice from the literature and experiments are that retention patches as large as 0.5 ha and 10-m-wide buffers to watercourses are not enough to maintain pre-harvest species composition but survival of forest species is still larger than on conventional clearcuts. Deadwood on clearcuts may present important habitats to saproxylic species, including rare and red-listed ones and a prioritization of tree species per stand is recommended. We identify several important future research directions including switch of focus towards the landscape as well as the species population level. Surveys in parts of European Russia where retention has been unintentionally implemented already for a century would indicate possible future trajectories of biodiversity and their drivers in other regions of Northern Europe. A stronger link to ecological theory would help in study designs and in the formulation of predicted outcomes.

Keywords: Biodiversity, Buffer strips, Conservation, Deadwood, Experiments, Forestry, High stumps, Retention patches, Variable retention

Background

Retention forestry (variable retention, variable retention forestry, retention silvicultural system, retention system, see Beese et al. 2019) is a relatively new forestry management concept in which measures to promote biodiversity are integrated into harvest operations (Gustafsson et al. 2012). It has expanded globally during the last decades and is today practiced on several continents (Lindenmayer et al. 2019; Palik and D'Amato 2019; Martínez Pastur et al. 2019; Shorohova et al. 2019), most commonly at clearcutting operations but the approach is

equally justified at selective harvest (Gustafsson et al. 2019). Main actions are to leave individual living trees and tree groups (retention patches) and often also to maintain or create deadwood at site, with provisioning of biodiversity as a key important objective (Martínez Pastur et al. 2009; Lindenmayer et al. 2012). Other objectives include structural enrichment of the post-harvest stand, secured continuity of structures and processes, maintenance of connectivity at the landscape scale, minimization of off-site impacts, and improved esthetic perception (Gustafsson et al. 2012).

Retention approaches were introduced in Northern Europe already in the 1980s (Simonsson et al. 2015), but large-scale application did not start until the late 1990s, when national certification standards (FSC, PEFC) were launched, and

* Correspondence: lena.gustafsson@slu.se

¹Department of Ecology, Swedish University of Agricultural Sciences, P.O. Box 7044, SE-75007 Uppsala, Sweden

Full list of author information is available at the end of the article

certification schemes were widely implemented by a large share of forest owners (Gustafsson et al. 2010). Research on ecological effects of retention is extensive worldwide (Lindenmayer et al. 2012), and several large-scale experiments have been established in different parts of the globe (Gustafsson et al. 2012). Meta-analyses demonstrate benefits to several species groups (Rosenvald and Löhmus 2008; Mori and Kitagawa 2014; Fedrowitz et al. 2014). A summary of research in Finland, Norway, and Sweden was published about a decade ago and listed about 50 biodiversity-oriented articles, with a dominance of studies on beetles and deadwood (Gustafsson et al. 2010). Due to increasing amount of research and newly established experiments in Northern Europe, this overview needs to be expanded and updated, including a systematic approach to literature retrieval. Furthermore, a summary of research published in the Russian language, so far largely unavailable to most researchers, is urgent.

Level of retention varies greatly between countries and regions; Gustafsson et al. (2012) report a range from a few percent of the logged area up to almost 50%, in a global comparison. Data on current state suitable for comparisons between North European countries are lacking, but an average of 11% of the area of harvested stands plus about 10 trees with special habitat value were recently reported from Sweden (Skogsstyrelsen 2019; Sveriges Officiella Statistik 2019). In Finland up to about 2% of the total volume of the growing stock is retained in final felling operations (Kuuluvainen et al. 2019). An indication of large differences between countries is a compilation of prescriptions on buffer zone widths along lakes and watercourses on forestland in the Nordic-Baltic region (Ring et al. 2017), revealing a very large between-country variation. Here we present an overview of research on retention approaches in Northern Europe (Norway, Sweden, Finland, the Baltics states, and NW Russia) with special emphasis on aspects of biodiversity. The aim of our study is to (1) compile and synthesize research on retention forestry from the region, (2) identify important future research directions, and (3) draw conclusions for practice. We divide the research synthesis (aim 1) into three separate parts. First, we analyze systematically retrieved literature on retention forestry from Sweden, Finland, Norway, and the Baltic states, published in peer-reviewed scientific journals. A compilation is also made of so far largely unknown literature from European Russia, published in the Russian language. Finally, we describe and present results from large-scale, replicated retention experiments conducted in Finland and Sweden.

Literature on retention in Northern Europe excluding Russia

Systematic search

We made a search in Web of Science Core Collection, Biosis Citation Index, and CAB Abstracts at the library

of Swedish University of Agricultural Sciences on August 9, 2019, using systematic literature retrieval methodology (Collaboration for Environmental Evidence 2018). We used the following search string: (retention OR retain*) AND (forest*) AND (Sweden OR Finland OR Norway OR Estonia OR Latvia OR Lithuania OR Fennoscandia OR Nordic OR Scandinavia OR Baltic). We applied the following criteria for filtering of articles: (1) peer-reviewed scientific natural science article published in English, (2) focusing on forest, retention, biodiversity, terrestrial environments, and (3) study conducted in Sweden, Finland, Norway, Estonia, Latvia, or Lithuania. A total of 1760 articles were found, reduced to 271 after reading title, 162 after reading abstract, and 120 after reading the whole text. A total of 59 articles were added from other sources, and thus the final compilation embraced a total of 179 articles. Articles were classified into their main research direction, and knowledge was synthesized for topics covered by several studies (border zones, retention patches, high stumps, other types of deadwood, *Populus tremula*, cost-efficiency) (see Additional file 1: Appendix A).


Characterization of studies

Most of the about 180 studies were performed in Sweden (51%), followed by Finland (31%), Estonia (10%), Norway (5%), and Latvia (2%) while only one originated from Lithuania. Only 5 articles were published before 2001, while 57% were published during the period 2001–2010, and 40% after that. Beetles were the most common study organisms (39% of studies), followed by lichens and bryophytes (both 16%). Deadwood was in focus for most studies (34%) compared with about 25% for retention patches and solitary/dispersed trees, respectively (Fig. 1, see also Additional file 1: Appendix A).

Border zones along watercourses

About 15 studies have evaluated biodiversity effects of leaving border zones edging to watercourses, with research conducted in Sweden, Norway, and Finland. One review has been published, focusing on water quality and biodiversity (Kuglerová et al. 2014).

Several studies have been conducted in the same study area in boreal mid-Sweden with retained (10 m on each side) or logged border zones to streams, and with forest in nature reserves as control. A general finding from this study area is that buffer strips maintain species richness and abundance of several organism groups, including red-listed species, better than open, logged areas, but worse than unlogged forest. After 10–11 years, 60% of the pre-harvest bryophyte species were found in retained border zones but only 20% in clearcut borders (Hylander and Weibull 2012). Number of species and individuals of land snails showed a similar trend (Hylander et al. 2004).



In a follow-up on lichens, local extinction was time-lagged; a significant decrease in richness of red-listed/indicator and calicioid species in the retained buffer strips was found 17 years after logging but not after 3 years (Johansson et al. 2018). A chronosequence study from the same region in Sweden showed that recovery of bryophytes after logging of border zones might take a long time. Streamside borders logged 30–50 years ago had an impoverished bryophyte flora compared to borders in older forests (Dynesius and Hylander 2007).

In a Finnish study on moths, buffer strips were considerably wider, 30–70 m, and were compared with logged borders and reference forests. Number of species and individuals did not differ between buffers and control forests but were significantly lower in clearcut borders (Mönkkönen and Mutanen 2003). In another Finnish study, buffer strips of at least 30 m on both sides of streams were needed to maintain species composition of vascular plants but for mosses 15 m sufficed (Oldén et al.

2019). For birds in Norway (Hågvar et al. 2004), the number of species increased with buffer width, up to 30 m from the watercourse. Strips 11–20 m wide had the highest density of species and individuals per hectare.

Retention patches

About 30 scientific articles have been published on retention patches in Northern Europe with > 90% conducted in Sweden and Finland, and with insects, lichens, and bryophytes being the most commonly studied organism groups. The first study on retention in Northern Europe was published in 1994 (Esseen 1994), reporting tree-uprooting in Norway spruce *Picea abies* L. H. Karst-dominated retention patches of different sizes. In a follow-up study 18 years later main conclusions were that patches as large as 1 ha are affected by wind disturbance and that most trees fall within the first 5 years while later on mortality is only a few percent per year (Jönsson et al. 2007). Early, high tree mortality was also

found in an Estonian study monitoring > 3000 dispersed retention trees over a period of 6 years (Rosenvald et al. 2008). Norway spruce has the highest mortality while Scots pine *Pinus sylvestris* L. and birch *Betula* spp. are less sensitive (Hallinger et al. 2016), and the risk of wind-felling is larger on paludified than on mineral soil (Hautala and Vanha-Majamaa 2006).

Only a few studies have focused on biodiversity responses to the size of retention patches, up to 0.6 ha in size. In a Finnish study, spider communities changed less after logging in large than in small patches at paludified sites (Matveinen-Huju et al. 2009), and two lichen species, *Calicium parvum* Tibell and *Micarea globulosella* (Nyl.) Coppins, decreased more in small than large patches when re-inventoried after 6 years in Central Sweden (Perhans et al. 2009). Time-series with repeated species recordings in retention patches are overall uncommon. In Perhans et al. (2009), the number of records of species of conservation concern decreased 30% for bryophytes while only 1% for lichens. In another Swedish study, exact locations of red-listed bryophytes and lichens were determined before harvest, re-located 3–7 years post logging, and were compared with unlogged control forest (Rudolphi et al. 2014). About half of bryophyte occurrences on logs remained in retention patches as well as in control forests, while only 10% were re-discovered on open clearcuts. About half of the lichen occurrences on living trees in patches remained compared with 75% in controls. Rudolphi et al. (2014) is the only study so far to assess the impact of location of patches. These authors found that the survival of desiccation-sensitive species may increase if patches are located in shady positions; local extinction was lower in north-facing than in south-facing patches, and in patches located close to mature forest.

High stumps

Every year, at least one million high stumps (stumps cut at a height of 3–5 m) in final felling and thinning are set in Sweden, aiming to support deadwood-associated species. This measure started in Sweden in the late 1990s and is nowadays also applied in other parts of the Nordic region. Here, 26 studies on this type of deadwood are summarized, of which 24 were conducted in Sweden. Many of these reports are on spruce stumps sampled within a few summers after cutting (Schroeder et al. 1999; Abrahamsson and Lindbladh 2006; Hedgren 2007; Lindbladh and Abrahamsson 2008; Abrahamsson et al. 2008; Fossetøl and Sverdrup-Thygeson 2009). Most studies are on a small spatial scale (tree), but some general patterns can be identified. First, high stumps differ in species composition from ordinary harvesting stumps (Abrahamsson and Lindbladh 2006; Hedgren 2007; Hjältén et al. 2010) and logs (Jonsell and Weslien 2003; Gibb

et al. 2006; Hjältén et al. 2007, 2010; Fossetøl and Sverdrup-Thygeson 2009). Second, species composition of insects and fungi differs considerably between tree species (Jonsell et al. 2004; Lindhe et al. 2004, 2005; Lindhe and Lindelöw 2004; Lindbladh et al. 2007). Third, diameter has little effect on early succession spruce communities (Jonsell and Weslien 2003; Lindbladh et al. 2007; Abrahamsson et al. 2008) whereas diameter seems to affect communities of species associated with deciduous tree and late successions (Jonsell et al. 2004; Lindhe et al. 2004; Lindhe and Lindelöw 2004; Weslien et al. 2011). Fourth, and important from a conservation point of view, many red-listed species occupy aspen high stumps (Jonsell et al. 2004; Lindhe et al. 2005). At the landscape scale, two studies explore how high stumps benefit recruitment of beetles (Schroeder et al. 2006; Djupström et al. 2012), and three explore how landscape structure affect the beetle fauna of high stumps of spruce and birch (Lindbladh et al. 2007; Abrahamsson et al. 2009; Jonsell et al. 2019). The vast majority of studies deal with beetles, two are on solitary bees and wasps nesting in cavities of dead trees (Westerfelt et al. 2015, 2018), two focus on parasitoids of early successional beetles (Hilszczański et al. 2005; Hedgren 2007), and one is on polypore fungi (Lindhe et al. 2004). In contrast to much other deadwood that is retained during cutting, the time of tree death is always known for high stumps. High stumps also remain standing during a long time and can be followed for decades of decay. Thus, high stumps have initiated novel studies on ecological succession which have added to the understanding of how communities assemble in deadwood (Weslien et al. 2011; Jacobsen et al. 2015; Jonsell et al. 2019). The studies are summarized in Table 1 with respect to stump age and investigated spatial scale.

Other types of deadwood

How important is deadwood retention at final harvest for supplying habitats for saproxylic species? The answer can partly be found among the about 30 papers included in this section. Although the volumes of dead trees in clearcuts have increased during the last decades (Kruys et al. 2013), they still constitute a small part of the total volumes in closed forests in the surrounding landscape (Schroeder et al. 2006; Ekbohm et al. 2006). Consequently, short-term responses of recruitment among saproxylic species should be found among those species that avoid shaded habitats in the closed forest and instead prefer sun-exposed deadwood on clearcuts, mostly insects.

European aspen *Populus tremula* L. (see below) hosts the highest proportion of sun-promoted beetle species followed by common oak *Quercus robur* L. and birches *Betula pendula* Roth, *B. pubescens* Ehrh. (Lindhe and

Table 1 Studies on high stumps by age (years after cutting) and spatial scale. The same set of stumps may recur in several papers but with different age, and several spatial scales can be addressed in the same paper

Years after cutting	Tree	Stand	Landscape
< 2	Schroeder et al. 1999; Hilszczański et al. 2005; Lindhe et al. 2005; Gibb et al. 2006; Johansson et al. 2006; Abrahamsson and Lindblad 2006; Hjältén et al. 2007; Hedgren 2007; Fossetøl and Sverdrup-Thygeson 2009	Hilszczański et al. 2005; Johansson et al. 2006; Hjältén et al. 2007; Lindblad and Abrahamsson 2008	
3–6	Jonsell and Weslien 2003; Jonsell et al. 2004, 2005; Lindhe et al. 2004, 2005; Lindhe and Lindelöw 2004; Abrahamsson et al. 2008; Westerfelt et al. 2015	Lindblad et al. 2007; Lindblad and Abrahamsson 2008; Westerfelt et al. 2018	Schroeder et al. 2006; Lindblad et al. 2007
7–10	Lindhe et al. 2004, 2005; Lindhe and Lindelöw 2004; Westerfelt et al. 2015	Westerfelt et al. 2018	Schroeder et al. 2006; Abrahamsson et al. 2009; Jonsell et al. 2019
≥ 11	Weslien et al. 2011; Jacobsen et al. 2015; Westerfelt et al. 2015	Westerfelt et al. 2018	Djupström et al. 2012

Lindelöw 2004; Lindhe et al. 2005). Aspens retained in clearcuts are important habitats for sun-loving insect species (Martikainen 2001; Sverdrup-Thygeson and Ims 2002; Sahlin and Ranius 2009; Sverdrup-Thygeson and Birkemoe 2009; Ranius et al. 2011). Birch stems in clearcuts harbor different species than those of similar diameter and decay stage inside forests (Kaila et al. 1997). Even small diameter retained birch stems can be valuable in this respect (Rubene et al. 2014). Clearing around ancient oaks in spruce plantations attracts beetle species that breed in dead branches of these oaks (Widerberg Koch et al. 2012). Spruce is shade tolerant which applies also for many saproxylic insects colonizing spruce stems (Lindhe and Lindelöw 2004; Lindhe et al. 2005; Gibb et al. 2006; Johansson et al. 2006; Hjältén et al. 2007; Fossetøl and Sverdrup-Thygeson 2009; Djupström et al. 2012).

Solitary bees and wasps nest almost exclusively in sun-exposed habitats and thus retained standing dead trees are important nesting sites for some species (Westerfelt et al. 2015, 2018). For saproxylic fungi, the degree of sun-exposure seems to be of minor importance compared to decay stage and diameter (Lindhe et al. 2004; Junninen et al. 2007).

Modeling of long-term effects of green tree and deadwood retention shows that retention levels of today affect deadwood amounts and qualities in future forest landscapes (Ranius & Kindvall 2004, Roberge et al. 2015). Some studies also show detrimental effects of forestry operations on deadwood amounts, particularly soil scarification (Hautala et al. 2004) and fuel wood extraction (Rudolphi and Gustafsson 2005).

One important question is whether one should retain deadwood in many small or a few large patches. Several studies indicate that species occupancy often increases with increasing substrate density (Komonen et al. 2000; Ranius 2002; Sahlin and Schroeder 2010; Rubene et al. 2014; Ranius et al. 2017). This would call for large habitat patches meaning retention of many stems of the same tree species per clearcut.

European aspen *Populus tremula*

European aspen is a keystone species with particular importance for biodiversity in northern Fennoscandia (Kouki et al. 2004). It is the most studied individual retention tree species with a total of about 25 articles. Several studies show that dead aspens in open, sun-exposed conditions are important for many species both for survival of pre-harvest species and for colonization. Fallen retention aspens have a more diverse flora of saproxylic fungi compared to other tree species (Runnel et al. 2013). Compared to mature forest, dead aspens on clearcuts host more polypore species (Junninen et al. 2007), and as about equally many beetle species (Martikainen 2001; Sverdrup-Thygeson and Ims 2002). Many beetle species on man-made high stumps of aspen thrive in sunny or semi-open conditions (Lindhe et al. 2005), and such stumps provide as important habitat as natural snags (Jonsell et al. 2004). Epiphytic lichens adapted to closed canopy conditions may survive on solitary live aspens, and with time, more light-tolerant species will colonize (Löhmus and Löhmus 2010; Lundström et al. 2013). The red-listed lichen *Lobaria pulmonaria* (L.) Hoffm. can survive and grow on retained green aspens, particularly on their northern side (Gustafsson et al. 2013). Epiphytic bryophytes are more sensitive than epiphytic lichens (Lundström et al. 2013; Oldén et al. 2014), but *Orthotrichum gymnostomum* Bruch ex Brid. is an example of a rare moss species that thrives on solitary retention aspens (Rudolphi et al. 2014). Volume of aspen and number of epiphytic lichens usually correlate positively (Hedenås and Ericson 2000; Lundström et al. 2013). However, many small aspen trees can host as many species as a few large trees, counted per bark area (Schei et al. 2013).

Cost efficiency

The relationship between lost wood production and biodiversity has been studied in about 10 articles. A Swedish project found that leaving dead trees and producing

high stumps was a cost-efficient way of creating habitats for red-listed insects and cryptogams (Ranius et al. 2005; Jonsson et al. 2006, Jonsson et al., 2010). Extended rotation time was, however, as efficient as artificial creation of deadwood. Leaving birch and aspen was more cost-efficient than leaving spruce and pine in southern Sweden, while conifers were more cost-efficient in the north. The cost-benefit ratio may be lower if retention patches are selected based on combined information on species content and timber volume (Perhans et al. 2011). Maintaining woodland key habitats (important sites for biodiversity) may, however, be the most cost-efficient conservation measure, based on the number of rare species, but retention patches are also cost-efficient due to their usually low timber volumes (Wikberg et al. 2009). By selecting small and crooked aspens with black bark as retention trees, many lichens can be preserved with low cost (Perhans et al. 2014). Retention of solitary pine trees in clearcuts reduces the growth of surrounding trees by a few percent, but losses can be compensated for by increased growth of the retained pines (Jakobsson and Elfving 2004; Elfving and Jakobsson 2006). A Finnish simulation study including different conservation scenarios and using habitat for deadwood-associated species as a proxy for biodiversity, concluded that compared to larger set-aside areas, small retention patches (0.01 ha) are an ineffective way to spend conservation funding (Mönkkönen et al. 2011).

Research in European Russia

Russian literature was not included in the systematic search since such studies have rarely been published in peer-reviewed, English-language scientific journals. Thus, we give a short overview of publications on retention trees and patches that have been published in the Russian language. In NW Russia (reaching from Finland and the Baltic states to the Ural mountains), forest patches, individual trees, and deadwood have been commonly retained at harvesting areas from the 1930s to the 1960s, mainly due to incomplete clear felling (Shorohova et al. 2019). During the last 20 years, the role of such unintentional retention has been documented in some monitoring reports and observational case studies. In the Republic of Karelia, many rare and threatened tree-associated birds, small mammals, wood-inhabiting fungi, lichens, and bryophytes occupy uncut forest patches surrounded by vast clearcuts or young forests (Kravchenko 1999; Gromtsev et al. 2010; Ivanter and Kurhinen 2016). Forest buffers on wet sites along rivers and around springs are of special importance for biodiversity and ecological integrity (Kravchenko 1999; Markovsky and Iljina 2014). Patches of undergrowth, seed tree patches and uncut patches of trees with non-merchantable wood may act as refugia for late-successional forest lichens, vascular plants,

and bryophytes after felling (Kravchenko et al. 2004; Kryshen 2006; Fadeeva and Kravchenko 2018).

Studies on contemporary, intentional retention are scarce. Inventories in the Pskov model forest (southwesternmost part of the region) demonstrate an association of many rare and threatened species of wood-inhabiting fungi (Kotkova 2010), lichens (Glushkovskaya 2010) and bryophytes (Kushnevskaia 2010) to retention elements and tree groups. In a study covering seven consecutive years following harvest, the total number of species and number of threatened species of invertebrates, vascular plants, bryophytes, polypores, and lichens were higher in areas where deadwood and special biotopes (such as paludified patches and creeks) had been retained than on open clearcuts (Romanyuk et al. 2009). Compilations made by species experts also show that many red-listed and indicator species are associated with certain structural elements and biotopes (Andersson et al. 2009; Markovsky and Iljina 2014).

Retention experiments in Finland and Sweden

Ecological responses to retention approaches have been studied experimentally in Finland since 1995, following the establishment of the “Biodiversity in regeneration of managed forests” (MONTA) experiment in southern Finnish spruce forests (Table 2 and Figs. 2 and 3). In Sweden, the only retention experiment so far, Effaråsen, was established in 2012.

Following the MONTA experiment (W and E Finland; Fig. 2), four other experiments have been established in Finland. The small retention-tree groups in MONTA (0.01–0.09 ha, 20–30 trees per group) were not able to maintain the pre-treatment forest species and communities of, e.g., understory vegetation (especially mosses and liverworts) and ground beetles (Jalonen and Vanha-Majamaa 2001; Koivula 2002a). Following this observation, RETREE was designed in 1997 to determine whether larger, up to 0.6 ha in size, retention-tree groups can maintain the pre-treatment (forest and moist habitat) species better than the small ones (Hautala et al. 2011). In 2000, the FIRE experiment was established in Eastern Finnish pine-dominated forests to study variable retention with and without prescribed burning. In 2001, the EVO experiment was established in southern Finnish spruce forests with a somewhat different study design and aims, the focus being on restoration measures (downed deadwood and burning). Here, an additional aspect was the comparison between upland and paludified biotopes within each stand, allowing an assessment of interactions between management operations and moisture. The most recent Finnish experiment is the “Forest management inspired by natural disturbance dynamics” DISTDYN (W and E Finland; Fig. 2), established in

Table 2 Experiments on retention forestry in Finland and Sweden. The Finnish experiments are located in the middle boreal biome and the Swedish experiment in the north boreal biome (Ahti et al. 1968). The Finnish experiments are ordered according to establishment year. For location of experiments, see Fig. 2, and for photos from a selection of experiments, see Fig. 3. *Year* establishment year, *Age* stand age at establishment (year), *NS* Norway spruce *Picea abies*, *SP* Scots pine *Pinus sylvestris*. Blocks, size = number of blocks or stands, and their size ranges. *LUKE* Natural Resources Institute Finland, *Metsähallitus* Finnish forest and park services, *Metsäteho* Metsäteho Ltd., a research and development company by Finnish forest industry organizations, *UEF* University of Eastern Finland, *UH* University of Helsinki, *Skogforsk* Forestry Research Institute of Sweden. For explanation of Finnish experiment names, see the section “Abbreviations” in the text


Experiments	Characteristics							
	Year	Age, tree species	Blocks, size	Treatments	Response variables	Responsible	Information	
MONTA (East and West) (Finland)	1995	90, NS	48, 1–2 ha	Clear felling, retention felling (up to 0.02 ha patches), gap felling, selection felling	Wood-inhabiting fungi, vascular plants, lichens, mosses, liverworts, beetles, spiders, ants, land snails, timber volume, sapling establishment, coarse woody debris	LUKE, Metsäteho, UH	Jalonen and Vanha-Majamaa 2001	
RETFREE (Finland)	1997	90, NS	11, 1–3 ha	Retention of 0.1–0.6 ha patches in clear-felled stands	Vascular plants, lichens, mosses, liverworts, beetles, spiders, land snails, timber volume, sapling establishment, coarse woody debris	LUKE, UH	Matveinen-Huju et al. 2006	
FIRE (Finland)	2000	150, SP	24, 3–5 ha	Retention of different intensity (0, 10, or 50 m ³ /ha), burning	Polypore and other fungi, beetles, flat bugs, pollinators, small mammals, ungulates, lichens, tree monitoring, tree regeneration, dendrochronology	UEF	Hyvärinen et al. 2005	
EVO (Finland)	2001	90, NS	24, 1–3 ha	Down wood retention (5, 30, or 60 m ³ /ha), standing tree retention (50 m ³ /ha), burning	Wood-inhabiting fungi, vascular plants, lichens, mosses, liverworts, beetles, timber volume, sapling establishment, coarse woody debris	LUKE, UH	Vanha-Majamaa et al. 2007	
DISTDYN (East and West) (Finland)	2009	90, NS/SP	12, 200–300 ha	Selection felling, gap felling, clear felling, deadwood addition. 5–20% retention volume	Wood-inhabiting fungi, vascular plants (some lichens and mosses), beetles, timber volume, sapling establishment	LUKE, Metsähallitus, UEF, UH	Koivula et al. 2014	
Effaråsen (Sweden)	2012	120, SP	24, 5–14 ha	Retention gradient (3–100% volume) and burning	Wood-inhabiting fungi, mycorrhiza fungi, epixylic lichens, beetles, timber volume, logging performance and opportunity costs, carbon stock, albedo	Skogforsk	Djupström and Weslien 2019	

2009, expanding the scope from stand-level comparisons to landscape-level assessments (Koivula et al. 2014).

Publications from the Finnish experiments so far cover 10–20 post-harvest years. Their main findings can be summarized as follows: (1) selection felling removing 33% of stand volume (dispersed retention), supports most of the common forest species, including ground beetles, spiders, soil fauna, and understory vegetation (Koivula 2002b; Matveinen-Huju and Koivula 2008; Siira-Pietikainen and Haimi 2009; Vanha-Majamaa et al. 2017), (2) retention felling with 7% retained volume in tree groups (aggregated retention) of about 0.05 ha mitigate some faunal and floral alterations (Jalonen and Vanha-Majamaa 2001; Koivula 2002a; Siira-Pietikainen and Haimi 2009; Koivula et al. 2019); however, even considerably larger retention patches (0.6 ha) do not maintain species composition of mosses, liverworts, or lichens (Hautala et al. 2011; Hämäläinen et al. 2016) and species composition of more common species is also altered (Matveinen-Huju et al. 2006, 2009), (3) gap felling with up to 50% of the stand clearcut as three 30 × 40 m gaps mitigate effects of clear-felling via their retained

parts, but also their cleared parts are less altered than 1–2-ha clearcuts (Koivula and Niemelä 2003; Matveinen-Huju and Koivula 2008), (4) as partly reflected above, a common finding in retention experiments is that the higher the retention, the better it mitigates the effects of clearfelling (Hyvärinen et al. 2005; Martikainen et al. 2006; Toivanen and Kotiaho 2010; Toivanen et al. 2014; Johnson et al. 2014; Rodríguez and Kouki, 2015; Heikkala et al. 2016; Vanha-Majamaa et al. 2017; Jokela et al. 2019; Suominen et al. 2019), (5) restoration efforts, such as addition of deadwood and/or prescribed burning, support rare and red-listed species (Hyvärinen et al. 2006; Toivanen and Kotiaho 2007; Suominen et al. 2015, 2019; Heikkala et al. 2017), but some species groups containing many red-listed species are negatively affected, such as lichens (Hämäläinen et al. 2014; Hämäläinen 2016), and (6) closed-forest species benefit from moist patches and high micro-habitat variation at harvested sites (Koivula 2002a; Toivanen et al. 2014; Koivula and Vanha-Majamaa 2020, in press).

For the Swedish Effaråsen experiment, baseline data and modelling studies have been published. Logging



time (minutes per m^3 wood), opportunity costs, and deadwood diversity increased with an increasing retention level (Santaniello et al. 2016). Baseline data show that old hard pine wood is a valuable habitat for epixylic lichens (Santaniello et al. 2017a). Modelling approaches include wood production, deadwood accumulation, and climate-related issues on carbon stocks and albedo (Santaniello et al. 2017b; Cherubini et al. 2018).

Reflections on retention forestry research in Northern Europe

Our analysis of peer-reviewed publications, overview of grey literature from NW Russia, and compilation of findings from experiments in Finland and Sweden demonstrate that substantial research efforts have been made in Northern Europe to better understand the relationship between biodiversity and the retention forestry system. Our findings are valid also for other boreal and temperate forest regions, since the retention forestry system has notable similarities in application in different biomes (Gustafsson et al. 2012). The number of articles (almost 200) was highest in the period 2006–2010 (about 60 studies) but the interest continues, and the last 3 years more than 20 papers have been published. Already almost a decade ago, Lindenmayer et al. (2012) reported

about 500 studies on retention approaches and biodiversity globally, with publications from Northern Europe comprising about 20%. Diverse aspects have been studied in the Nordic and Baltic countries but the focus on deadwood is particularly prominent; almost 35% of all studies have assessed this component. High stumps have been the target of more than 25 studies, which might be surprising but there is a special interest in this type of deadwood since high stumps are associated with certification. In Sweden for instance, both FSC and PEFC require a creation of at least three high stumps per ha at each harvest occasion. Another characteristic of studies from Northern Europe is the large emphasis on invertebrates, bryophytes, lichens, and fungi (focus in 90% of studies), and less so on the better-known taxonomic groups of vascular plants and vertebrates. Remarkably, no study has been done on mammals. Most studies have been performed in Sweden and Finland (80% of all publications), and also all large-scale, replicated experiments have been established in these countries, with a striking concentration to Finland (5 out of 6 experiments). The Russian literature corroborates the peer-reviewed publications and demonstrates the importance of (intentionally or unintentionally) retained patches and buffer zones as refugia for organisms. A similar focus on lesser known



Fig. 3 Examples of retention experiments in Finland and Sweden. **a** Gap felling with 5% group retention inside a gap, spruce forest, third post-harvest summer. DISTDYN project in Finland. Photo: Matti Koivula. **b** In the Swedish experiment, Effaråsen, Scots pine stands have been harvested with retention levels varying from 3 to 100%. At each level the retention included equal proportions of intact green trees, girdled trees, high-cut trees, and felled trees. In the picture, a level of 30% retention is shown. Photo Line Djupström. **c** Retention tree group inside a clear-felled area. In the RETREE project, varying sized retention tree groups, 0.1–0.6 ha were left in a harvested area. Photo: Erkki Oksanen/LUKE. **d**. Naturally regenerated gap in the MONTA experiment 14 years after harvesting. Photo: Erkki Oksanen/LUKE. For more information on the experiments, see Table 2, and for their location, see Fig. 2.

organisms (bryophytes, lichens, fungi) is also evident in studies from this part of Northern Europe. The overview on large-scale, replicated retention experiments given here supplements earlier descriptions of experiments performed around the globe (Gustafsson et al. 2012). Results from the experiments are clear; substantial amounts of retention are needed to counteract negative effects of logging on species associated with old forest, agreeing with meta-analyses on retention forestry and biodiversity (e.g., Fedrowitz et al. 2014). In selection felling as well as gap felling, > 50% of stand volume needs to be retained in order to maintain common forest species, and retention patches need to be more than half a ha in size, and probably larger, to preserve old-forest species communities. Important mechanisms for the long-term survival of old-forest species include moist microclimate and availability of various microhabitats. Insights into restoration efforts have also been gained from the experiments; many saproxylic species benefit from addition of deadwood and prescribed burning, actions compatible with retention forestry.

Future research

Continued research on retention approaches is urgent in Northern Europe, to evaluate current practices but also

to provide an improved knowledge base necessary for future development of forest management. Some research directions associated with biodiversity stand out as especially important. First, there is a need for upscaling to the landscape level, since many species depend on large-scale patterns and processes and to date almost all surveys and experiments have been done at the stand level. Second, there is a lack of population-oriented studies, since current literature is largely dominated by assessments of species richness and composition, measures that are difficult to evaluate. As management modifications should support species that need conservation efforts, i.e., rare and threatened species, their responses are particularly relevant. Third, the European part of Russia offers a unique opportunity to evaluate long-term effects of retention. In this region large areas have been logged since 1910 with mainly unintentionally retained single trees and groups of trees (Shorohova et al. 2019). Surveys of species as well as forest and landscape structures would indicate the future composition of biodiversity and its drivers in landscapes in which retention has been more recently introduced, like in other parts of Northern Europe. Fourth, although most studies to date have attempted to answer applied questions, we

recommend stronger links to ecological theory. Such would aid in study designs and in the formulation of predictions of outcomes, and strengthen the interpretation of results. Fifth, maintenance of current experiments is essential, and implementation of cross-country monitoring programs would greatly aid in the evaluation of benefits to biodiversity of current management models. Finally, our review only includes natural science literature but more social science studies and studies in humanities are also needed in order to reach a more complete understanding of the complex retention forestry system.

Conclusions for practice

The large amount of research from Northern Europe enables some broad conclusions for practical forest management. First, the importance to biodiversity associated with old forests and elements herein increases with retention level, i.e., the more that is retained, the better for forest plants, animals and fungi. Although a general finding from the Finnish experiments is that some taxa are negatively affected even in retention patches as large as 0.6 ha, studies also show that some old-growth-forest species, including many of special conservation concern, may survive in smaller retention patches and on dispersed trees. Thus, as much trees as possible should be retained and size of patches should be maximized but even leaving small amounts is better than traditional clearcutting with removal of all trees. Second, buffer strips need to be few tens of meters on each side of a stream to maintain pre-logging richness and composition of various taxa. Compared to logged buffers, several species including certain red-listed ones nevertheless may survive in narrower strips, i.e., the conclusion is the same as for retention patches: leaving a little is better than leaving nothing at all. Third, retention and creation of sun-exposed deadwood in the form of logs, snags, and high stumps on clearcuts should be encouraged since they form important habitats for saproxylic taxa such as certain beetles and fungi, including some rare and red-listed species. Fourth, assuming a given volume to be retained, the research indicates that concentration of deadwood of one tree species per site is more efficient than leaving deadwood of a diversity of tree species. Fifth, a conclusion put forward in many publications, with which we agree, is that retention forestry is not a panacea for biodiversity but instead one way to mitigate negative effects of clearcutting. For some forest species, large forest reserves are necessary for their long-term survival.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13717-019-0208-2>.

Additional file 1: Supplementary material. 3 Appendix A literature on retention (Excel file with 4 worksheets).

Abbreviations

DISTDYN (W and E): A large-scale and long-term research and development project "Forest management inspired by natural disturbance dynamics"; EVO: Experimental research project established in Evo to study the biodiversity effects of variable retention, burning and deadwood manipulation; FIRE: An experimental project "Boreal forest fire experiment," established in eastern Finland to study the biodiversity effects of variable retention and burning; MONTA (W and E): Experimental research project "Biodiversity in the regeneration of managed forest"; RETREE: Experimental research project to study the effects of variable retention tree group size on biodiversity

Acknowledgements

Not applicable

Authors' contributions

LG was the lead author in writing, made the systematic search and classification of literature, edited the manuscript, and prepared the figures. MH wrote the sections on aspen and cost-efficiency. ES wrote the section on the Russian literature. IVM and MK wrote the section on the Finnish experiments. JW wrote the sections on high stumps and deadwood, and the text on the Swedish experiment Effaråsen. All authors read and approved the final manuscript.

Funding

The State Research Programme of the Forest Research Institute of the Karelian Research Centre, Russian Academy of Sciences funded the work by ES.

Availability of data and materials

Not applicable

Ethics approval and consent to participate

Not applicable

Consent for publication

The authors consent to publish the data included in this paper.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Ecology, Swedish University of Agricultural Sciences, P.O. Box 7044, SE-75007 Uppsala, Sweden. ²Silvinformation AB, Bergauddsvägen 22B, SE-39359 Kalmar, Sweden. ³School of Forest Sciences, University of Eastern Finland, P.O. Box 111, FI-80101 Joensuu, Finland. ⁴Forest Research Institute of the Karelian Research Centre, Russian Academy of Sciences, Pushkinskaya str. 11, Petrozavodsk 185910, Russia. ⁵Natural Resources Institute Finland (Luke), Latokartanonkaari 9, FI-00790 Helsinki, Finland. ⁶Saint-Petersburg State Forest Technical University, Institut'sky str. 5, Saint Petersburg 194021, Russia. ⁷The Forestry Research Institute of Sweden (Skogforsk), SE-75183 Uppsala, Sweden.

Received: 8 October 2019 Accepted: 12 December 2019

Published online: 17 January 2020

References

- Abrahamsson M, Jonsell M, Niklasson M, Lindbladh M (2009) Saproxylic beetle assemblages in artificially created high-stumps of spruce (*Picea abies*) and birch (*Betula pendula/pubescens*) - does the surrounding landscape matter? *Insect Conserv Divers* 2:284–294
- Abrahamsson M, Lindbladh M (2006) A comparison of saproxylic beetle occurrence between man-made high- and low-stumps of spruce (*Picea abies*). *For Ecol Manag* 226:230–237
- Abrahamsson M, Lindbladh M, Rönnerberg J (2008) Influence of butt rot on beetle diversity in artificially created high-stumps of Norway spruce. *For Ecol Manag* 255:3396–3403
- Ahti T, Hämet-Ahti L, Jalas J (1968) Vegetation zones and their sections in north-western Europe. *Ann Zool Fenn* 5:169–211
- Andersson L, Alekseeva N, Kuznetsova E (2009) Survey of biologically valuable forests in the northwestern European Russia. Volume 2. Guidebook for

- identification of species in the forest stand level of survey. StP, Pobeda Publ. 238 p. (in Russian)
- Beese WJ, Deal J, Dunsworth BG, Mitchell SJ, Philpott TJ (2019) Two decades of variable retention in British Columbia: a review of its implementation and effectiveness for biodiversity conservation. *Ecol Process* 8:33
- Cherubini F, Santaniello F, Hu X, Sonesson J, Hamner Strømman A, Weslien J, Djupström LB, Ranius T (2018) Climate impacts of retention forestry in a Swedish boreal pine forest. *J Land Use Sci* 13:301–318
- Collaboration for Environmental Evidence. 2018. Guidelines and Standards for Evidence synthesis in Environmental Management. Version 5.0 (AS Pullin, GK Frampton, B Livoreil & G Petrokofsky, Eds)
- Djupström L, Weslien J (2019) Forest management and biological conservation in old pine forests A long-term field experiment at Effaråsen in Sweden Part 1 2012–2018. Skogforsk, Uppsala
- Djupström LB, Weslien J, ten Hoopen J, Schroeder LM (2012) Restoration of habitats for a threatened saproxylic beetle species in a boreal landscape by retaining dead wood on clear-cuts. *Biol Conserv* 155:44–49
- Dynesius M, Hylander K (2007) Resilience of bryophyte communities to clear-cutting of boreal stream-side forests. *Biol Conserv* 135:423–434
- Ekbohm B, Schroeder LM, Larsson S (2006) Stand specific occurrence of coarse woody debris in a managed boreal forest landscape in central Sweden. *For Ecol Manag* 221:2–12
- Elfving B, Jakobsson R (2006) Effects of retained trees on tree growth and field vegetation in *Pinus sylvestris* stands in Sweden. *Scand J For Res* 21:29–36
- Esseen PA (1994) Tree mortality patterns after experimental fragmentation of an old-growth conifer forest. *Biol Conserv* 68:19–28
- Fadeeva M, Kravchenko A (2018) Lichens of clear cut areas in the southern Karelian forests. In: Fungal communities in forest ecosystems. Vol. 5. Moscow – Petrozavodsk. Pp: 122–135. (in Russian)
- Fedrowitz K, Koricheva J, Baker SC, Lindenmayer DB, Palik B, Rosensvald R, Beese W, Franklin JF, Kouki J, Macdonald E, Messier C, Sverdrup-Thygeson A, Gustafsson L (2014) Can retention forestry help conserve biodiversity? A meta-analysis. *J Appl Ecol* 51:1669–1679
- Fossestøl KO, Sverdrup-Thygeson A (2009) Saproxylic beetles in high stumps and residual downed wood on clear-cuts and in forest edges. *Scand J For Res* 24:403–416
- Gibb H, Pettersson RB, Hjältén J, Hilszczański J, Ball JP, Johansson T, Atlegrim O, Danell K (2006) Conservation-oriented forestry and early successional saproxylic beetles: Responses of functional groups to manipulated dead wood substrates. *Biol Conserv* 129:437–450
- Glushkovskaya N (2010) Lichenized fungi of Pskov model forest. In: The results of biodiversity survey in Pskov model forest. Ed. by Zagidullina AT and Romanyuk BD. Pp: 39–44 (in Russian)
- Gromtsev A, Kravchenko A, Kurhinen Y, Sazonov S (2010) Dynamics of forest communities, flora and fauna in the European taiga under natural conditions and after anthropogenic influence: an attempt of research and generalization. *Proc. Karelian Res. Centre RAS* 1:16–33 (in Russian)
- Gustafsson L, Baker SC, Bauhus J, Beese WJ, Brodie A, Kouki J, Lindenmayer DB, Löhmus A, Martínez Pastur G, Messier C, Neyland M, Palik B, Sverdrup-Thygeson A, Volney JA, Wayne A, Franklin JF (2012) Retention forestry to maintain multifunctional forests: A World Perspective. *BioScience* 62:633–645
- Gustafsson L, Bauhus J, Asbeck T, Augustynczik ALD, Basile M, Frey J, Gutzat F, Hanewinkel M, Helbach J, Jonker M, Knuff A, Messier C, Penner J, Pyttel P, Reif A, Storch F, Winiger N, Winke G, Yousefpour R, Storch I (2019) Retention as an integrated biodiversity conservation approach for continuous-cover forestry in Europe. *Ambio* 49:85–97
- Gustafsson L, Fedrowitz K, Hazell P (2013) Survival and vitality of a macrolichen 14 years after transplantation on aspen trees retained at clearcutting. *For Ecol Manag* 291:436–441
- Gustafsson L, Kouki J, Sverdrup-Thygeson A (2010) Tree retention as a conservation measure in clear-cut forests of northern Europe: a review of ecological consequences. *Scand J For Res* 25:295–308
- Håggvar S, Nygaard P, Bækken BT (2004) Retention of forest strips for bird-life adjacent to water and bogs in Norway: Effect of different widths and habitat variables. *Scand J For Res* 19:452–465
- Hallinger M, Johansson V, Schmalholz M, Sjöberg S, Ranius T (2016) Factors driving tree mortality in retained forest fragments. *For Ecol Manag* 368:163–172
- Hämäläinen A (2016) Retention forestry and intensified biomass harvest: epiphytic lichen assemblages under opposing ecological effects in pine-dominated boreal forests. Diss For 218. School of Forest Sciences Faculty of Science and Forestry, University of Eastern Finland. Joensuu.
- Hämäläinen A, Hujo M, Heikkala O, Junninen K, Kouki J (2016) Retention tree characteristics have major influence on the post-harvest tree mortality and availability of coarse woody debris in clear-cut areas. *For Ecol Manag* 369:66–73
- Hämäläinen A, Kouki J, Löhmus P (2014) The value of retained Scots pines and their dead wood legacies for lichen diversity in clear-cut forests: The effects of retention level and prescribed burning. *For Ecol Manag* 324:89–100
- Hautala H, Jalonen J, Laaka-Lindberg S, Vanha-Majamaa I (2004) Impacts of retention felling on coarse woody debris (CWD) in mature boreal spruce forests in Finland. *Biodivers Conserv* 13:1541–1554
- Hautala H, Laaka-Lindberg S, Vanha-Majamaa I (2011) Effects of retention felling on epixylic species in boreal spruce forests in southern Finland. *Restor Ecol* 19:418–429
- Hautala H, Vanha-Majamaa I (2006) Immediate tree uprooting after retention-felling in a coniferous boreal forest in Fennoscandia. *Can J For Res* 36:3167–3172
- Hedenäs H, Ericson L (2000) Epiphytic macrolichens as conservation indicators: successional sequence in *Populus tremula* stands. *Biol Conserv* 93:43–53
- Hedgren PO (2007) Early arriving saproxylic beetles (Coleoptera) and parasitoids (Hymenoptera) in low and high stumps of Norway spruce. *For Ecol Manag* 241:155–161
- Heikkala O, Martikainen P, Kouki J (2016) Decadal effects of emulating natural disturbances in forest management on saproxylic beetle assemblages. *Biol Conserv* 194:39–47
- Heikkala O, Martikainen P, Kouki J (2017) Prescribed burning is an effective and quick method to conserve rare pyrophilous forest-dwelling flat bugs. *Insect Conserv Divers* 10:32–41
- Hilszczański J, Gibb H, Hjältén J, Atlegrim O, Johansson T, Pettersson RB, Ball JP, Danell K (2005) Parasitoids (Hymenoptera, Ichneumonidae) of saproxylic beetles are affected by forest successional stage and dead wood characteristics in boreal spruce forest. *Biol Conserv* 126:456–464
- Hjältén J, Johansson T, Alinvi O, Danell K, Ball JP, Pettersson R, Gibb H, Hilszczański J (2007) The importance of substrate type, shading and scorching for the attractiveness of dead wood to saproxylic beetles. *Basic Appl Ecol* 8:364–376
- Hjältén J, Stenbacka F, Andersson J (2010) Saproxylic beetle assemblages on low stumps, high stumps and logs: Implications for environmental effects of stump harvesting. *For Ecol Manag* 260:1149–1155
- Hylander K, Nilsson C, Gothner T (2004) Effects of buffer-strip retention and clearcutting on land snails in boreal riparian forests. *Conserv Biol* 18:1052–1062
- Hylander K, Weibull H (2012) Do time-lagged extinctions and colonizations change the interpretation of buffer strip effectiveness? - a study of riparian bryophytes in the first decade after logging. *J Appl Ecol* 49:1316–1324
- Hyvärinen E, Kouki J, Martikainen P, Lappalainen H (2005) Short-term effects of controlled burning and green-tree retention on beetle (Coleoptera) assemblages in managed boreal forests. *For Ecol Manag* 212:315–332
- Hyvärinen E, Kouki J, Martikainen P (2006) Fire and green-tree retention in conservation of red-listed and rare deadwood-dependent beetles in Finnish boreal forests. *Conserv Biol* 20:1711–1719
- Ivanter E, Kurhinen Y (2016) Impact of commercial forest logging on the boreal fauna communities (small mammals of the Eastern Fennoscandia as an example). *Bull Russ Acad Sci Biol Series* 4:412–421 (in Russian)
- Jacobsen RM, Birkemoe T, Sverdrup-Thygeson A (2015) Priority effects of early successional insects influence late successional fungi in dead wood. *Ecol Evol* 5:4896–4905
- Jakobsson R, Elfving B (2004) Development of an 80-year-old mixed stand with retained *Pinus sylvestris* in Northern Sweden. *For Ecol Manag* 194:249–258
- Jalonen J, Vanha-Majamaa I (2001) Immediate effects of four different felling methods on mature boreal spruce forest understorey vegetation in southern Finland. *For Ecol Manag* 146:25–34
- Johansson T, Gibb H, Hilszczański J, Pettersson RB, Hjältén J, Atlegrim O, Ball JP, Danell K (2006) Conservation-oriented manipulations of coarse woody debris affect its value as habitat for spruce-infesting bark and ambrosia beetles (Coleoptera: Scolytinae) in northern Sweden. *Can J For Res* 36:174–185
- Johansson V, Wikström C-J, Hylander K (2018) Time-lagged lichen extinction in retained buffer strips 16.5 years after clear-cutting. *Biol Conserv* 225:53–65
- Johnson S, Strebom J, Kouki J (2014) Low levels of tree retention do not mitigate the effects of clearcutting on ground vegetation dynamics. *For Ecol Manag* 330:67–74
- Jokela J, Siitonen J, Koivula M (2019) Short-term effects of selection, gap, patch and clear cutting on the beetle fauna in boreal spruce-dominated forests. *For Ecol Manag* 446:29–37

- Jonsell M, Abrahamsson M, Widenfalk L, Lindbladh M (2019) Increasing influence of the surrounding landscape on saproxylic beetle communities over 10 years succession in dead wood. *For Ecol Manag* 440:267–284
- Jonsell M, Nitterus K, Stighäll K (2004) Saproxylic beetles in natural and man-made deciduous high stumps retained for conservation. *Biol Conserv* 118:163–173
- Jonsell M, Schroeder M, Weslien J (2005) Saproxylic beetles in high stumps of spruce: Fungal flora important for determining the species composition. *Scand J For Res* 20:54–62
- Jonsell M, Weslien J (2003) Felled or standing retained wood - it makes a difference for saproxylic beetles. *For Ecol Manag* 175:425–435
- Jonsson M, Ranius T, Ekvall H, Bostedt G (2010) Cost-effectiveness of silvicultural measures to increase substrate availability for wood-dwelling species: A comparison among boreal tree species. *Scand J For Res* 25:46–60
- Jonsson M, Ranius T, Ekvall H, Bostedt G, Dahlberg A, Ehnström B, Nordén B, Stokland JN (2006) Cost-effectiveness of silvicultural measures to increase substrate availability for red-listed wood-living organisms in Norway spruce forests. *Biol Conserv* 127:443–462
- Jönsson MT, Fraver S, Jonsson BG, Dynesius M, Rydgård M, Esseen PA (2007) Eighteen years of tree mortality and structural change in an experimentally fragmented Norway spruce forest. *For Ecol Manag* 242:306–313
- Junninen K, Penttilä R, Martikainen P (2007) Fallen retention aspen trees on clear-cuts can be important habitats for red-listed polypores: a case study in Finland. *Biodivers Conserv* 16:475–490
- Kaila L, Martikainen P, Punttila P (1997) Dead trees left in clear-cuts benefit saproxylic Coleoptera adapted to natural disturbances in boreal forests. *Biodivers Conserv* 6:1–18
- Koivula M (2002a) Alternative harvesting methods and boreal carabid beetles (Coleoptera, Carabidae). *For Ecol Manag* 167:103–121
- Koivula M (2002b) Boreal carabid-beetle (Coleoptera, Carabidae) assemblages in thinned uneven-aged and clear-cut spruce stands. *Ann Zool Fenn* 131–149
- Koivula M, Kuuluvainen T, Hallman E, Kouki J, Siitonen J, Valkonen S (2014) Forest management inspired by natural disturbance dynamics (DISTDYN) - a long-term research and development project in Finland. *Scand J For Res* 29:579–592
- Koivula M, Niemelä J (2003) Gap felling as a forest harvesting method in boreal forests: responses of carabid beetles (Coleoptera, Carabidae). *Ecography* 26:179–187
- Koivula M, Vanha-Majamaa I (2020) Experimental evidence on biodiversity impacts of variable retention forestry, prescribed burning and deadwood manipulation in Fennoscandia. *Ecol Process* (in press).
- Koivula MJ, Venn S, Hakola P, Niemelä J (2019) Responses of boreal ground beetles (Coleoptera, Carabidae) to different logging regimes ten years post harvest. *For Ecol Manag* 436:27–38
- Komonen A, Penttilä R, Lindgren M, Hanski I (2000) Forest fragmentation truncates a food chain based on an old-growth forest bracket fungus. *Oikos* 90:119–126
- Kotkova V (2010) Aphylophorales fungi in different forest habitats of Pskov model forest. In: *The results of biodiversity survey in Pskov model forest*. Ed. by Zagidullina AT and Romanyuk BD. Pp: 32–39 (in Russian)
- Kouki J, Arnold K, Martikainen P (2004) Long-term persistence of aspen - a key host for many threatened species - is endangered in old-growth conservation areas in Finland. *J Nat Conserv* 12:41–52
- Kravchenko A (1999) The role of uncut forest patches in preserving biodiversity in boreal forests after commercial harvesting. In: *Proc. Int. Conf. "Biological basis for study, use and preserve animals, plants and soils in the Eastern Fennoscandia"* Petrozavodsk, 6–10 Sept. P. 190–191 (in Russian)
- Kravchenko A, Gnatyuk E, Kryshen A (2004) Anthropogenic transformation of flora in the regions with intensive forest use. In: *Anthropogenic transformation of European boreal forest ecosystems: ecological, resource and economic aspects*. Proc. Int. Conf. Petrozavodsk, 23–25 of Dec. Pp. 82–93. (in Russian)
- Kruys N, Fridman J, Götmark F, Simonsson P, Gustafsson L (2013) Retaining trees for conservation at clearcutting has increased structural diversity in young Swedish production forests. *For Ecol Manag* 304:312–321
- Kryshen A (2006) Plant communities on the harvesting areas in Karelia. *Nauka publ, Moskow* (in Russian)
- Kuglerová L, Ågren A, Jansson R, Laudon H (2014) Towards optimizing riparian buffer zones: Ecological and biogeochemical implications for forest management. *For Ecol Manag* 334:74–84
- Kushnevskaia E (2010) Bryophytes of Pskov model forest. In: *The results of biodiversity survey in Pskov model forest*. Ed. by Zagidullina AT and Romanyuk BD. Pp: 44–49 (in Russian)
- Kuuluvainen T, Lindberg H, Vanha-Majamaa I, Keto-Tokoi P, Punttila P (2019) Low-level retention forestry, certification, and biodiversity: case Finland. *Ecol Process* 8:47
- Lindbladh M, Abrahamsson M (2008) Beetle diversity in high-stumps from Norway spruce thinnings. *Scand J For Res* 23:339–347
- Lindbladh M, Abrahamsson M, Seedre M, Jonsell M (2007) Saproxylic beetles in artificially created high-stumps of spruce and birch within and outside hotspot areas. *Biodivers Conserv* 16:3213–3226
- Lindenmayer D, Blair D, McBurney L (2019) Variable retention harvesting in Victoria's Mountain Ash (*Eucalyptus regnans*) forests (southeastern Australia). *Ecol Process* 8:2
- Lindenmayer DB, Franklin JF, Löhms A, Baker SC, Bauhus J, Beese W, Brodie A, Kiehl B, Kouki J, Martínez Pastur G, Messier C, Neyland M, Palik B, Sverdrup-Thygeson A, Volney J, Wayne A, Gustafsson L (2012) A major shift to the retention approach for forestry can help resolve some global forest sustainability issues. *Conserv Lett* 5:421–431
- Lindhe A, Åsenblad N, Toresson HG (2004) Cut logs and high stumps of spruce, birch, aspen and oak - nine years of saproxylic fungi succession. *Biol Conserv* 119:443–454
- Lindhe A, Lindelöw Å (2004) Cut high stumps of spruce, birch, aspen and oak as breeding substrates for saproxylic beetles. *For Ecol Manag* 203:1–20
- Lindhe A, Lindelöw Å, Åsenblad N (2005) Saproxylic beetles in standing dead wood density in relation to substrate sun-exposure and diameter. *Biodivers Conserv* 14:3033–3053
- Löhms A, Löhms P (2010) Epiphyte communities on the trunks of retention trees stabilise in 5 years after timber harvesting, but remain threatened due to tree loss. *Biol Conserv* 143:891–898
- Lundström J, Jonsson F, Perhans K, Gustafsson L (2013) Lichen species richness on retained aspens increases with time since clear-cutting. *For Ecol Manag* 293:49–56
- Markovsky A, Iljina O (2014) Methodological recommendations for preserving biological diversity during timber harvesting in the Vologda region. WWF. (in Russian)
- Martikainen P (2001) Conservation of threatened saproxylic beetles: significance of retained aspen *Populus tremula* on clearcut. *Ecol Bull* 49:205–218
- Martikainen P, Kouki J, Heikkala O (2006) The effects of green tree retention and subsequent prescribed burning on ground beetles (Coleoptera : Carabidae) in boreal pine-dominated forests. *Ecography* 29:659–670
- Martínez Pastur GJ, Lencinas MV, Cellini JM, Peri PL, Esteban RS (2009) Timber management with variable retention in *Nothofagus pumilio* forests of Southern Patagonia. *For Ecol Manag* 258:436–443
- Martínez Pastur GJ, Rosas YM, Manríquez MT, Toro Manríquez M, Huertas Herrera A, Miller JA, Cellini JM, Barrera MD, Peri PL, Lencinas MV (2019) Knowledge arising from long-term research of variable retention harvesting in Tierra del Fuego: where do we go from here? *Ecol Process* 8:24
- Matveinen-Huju K, Koivula M (2008) Effects of alternative harvesting methods on boreal forest spider assemblages. *Can J For Res* 38:782–794
- Matveinen-Huju K, Koivula M, Niemelä J, Rauha AM (2009) Short-term effects of retention felling at mire sites on boreal spiders and carabid beetles. *For Ecol Manag* 258:2388–2398
- Matveinen-Huju K, Niemelä J, Rita H, O'Hara RB (2006) Retention-tree groups in clear-cuts: Do they constitute "life-boats" for spiders and carabids? *For Ecol Manag* 230:119–135
- Mönkkönen M, Mutanen M (2003) Occurrence of moths in boreal forest corridors. *Conserv Biol* 17:468–475
- Mönkkönen M, Reunanen P, Kotiaho JS, Juutinen A, Tikkanen OP, Kouki J (2011) Cost-effective strategies to conserve boreal forest biodiversity and long-term landscape-level maintenance of habitats. *Eur J For Res* 130:717–727
- Mori AS, Kitagawa R (2014) Retention forestry as a major paradigm for safeguarding forest biodiversity in productive landscapes: A global meta-analysis. *Biol Conserv* 175:65–73
- Oldén A, Ovaskainen O, Kotiaho JS, Laaka-Lindberg S, Halme P (2014) Bryophyte species richness on retention aspens recovers in time but community structure does not. *PLoS One* 9(4):e93786
- Oldén A, Selonen VAO, Lehtonen E, Kotiaho JS (2019) The effect of buffer strip width and selective logging on streamside plant communities. *BMC Ecol* 19:9
- Palik BJ, D'Amato AW (2019) Variable retention harvesting in Great Lakes mixed-pine forests: emulating a natural model in managed ecosystems. *Ecol Process* 8:16
- Perhans K, Appelgren L, Jonsson F, Nordin U, Söderström B, Gustafsson L (2009) Retention patches as potential refugia for bryophytes and lichens in managed forest landscapes. *Biol Conserv* 142:1125–1133

- Perhans K, Glöde D, Gilbertsson J, Persson A, Gustafsson L (2011) Fine-scale conservation planning outside of reserves: Cost-effective selection of retention patches at final harvest. *Ecol Econ* 70:771–777
- Perhans K, Haight RG, Gustafsson L (2014) The value of information in conservation planning: Selecting retention trees for lichen conservation. *For Ecol Manag* 318:175–182
- Ranius T (2002) Influence of stand size and quality of tree hollows on saproxylic beetles in Sweden. *Biol Conserv* 103:85–91
- Ranius T, Ekvall H, Jonsson M, Bostedt G (2005) Cost-efficiency of measures to increase the amount of coarse woody debris in managed Norway spruce forests. *For Ecol Manag* 206:119–133
- Ranius T, Kindvall O (2004) Modelling the amount of coarse woody debris produced by the new biodiversity-oriented silvicultural practices in Sweden. *Biol Conserv* 119:51–59
- Ranius T, Martikainen P, Kouki J (2011) Colonisation of ephemeral forest habitats by specialised species: beetles and bugs associated with recently dead aspen wood. *Biodivers Conserv* 20:2903–2915
- Ranius T, Mestre L, Bouget C, Schroeder M (2017) Fragmentation effects on dead wood-dependent species associated with disturbed forest habitats: implications for stump harvesting. *Scand J For Res* 32:260–267
- Ring E, Johansson J, Sandström C, Bjarnadóttir B, Finér L, Libište Z, Lode E, Stupák I, Sætersdal M (2017) Mapping policies for surface water protection zones on forest land in the Nordic–Baltic region: Large differences in prescriptiveness and zone width. *Ambio* 46:878–893
- Roberge J-M, Lämås T, Lundmark T, Ranius T, Felton A, Nordin A (2015) Relative contributions of set-asides and tree retention to the long-term availability of key forest biodiversity structures at the landscape scale. *J Environ Manage* 154:284–292
- Rodríguez A, Kouki J (2015) Emulating natural disturbance in forest management enhances pollination services for dominant *Vaccinium* shrubs in boreal pine-dominated forests. *For Ecol Manag* 350:1–12
- Romanyuk B, Zagidullina A, Knize A (2009) Planning forestry on a nature conservation basis WWF. *Forest, Pskov Model* (in Russian)
- Rosenvald R, Löhms A (2008) For what, when, and where is green-tree retention better than clear-cutting? A review of the biodiversity aspects. *For Ecol Manag* 255:1–15
- Rosenvald R, Löhms A, Kiviste A (2008) Preadaptation and spatial effects on retention-tree survival in cut areas in Estonia. *Can J For Res* 38:2616–2625
- Rubene D, Wikars LO, Ranius T (2014) Importance of high quality early-successional habitats in managed forest landscapes to rare beetle species. *Biodivers Conserv* 23:449–466
- Rudolphi J, Gustafsson L (2005) Effects of forest-fuel harvesting on the amount of deadwood on clear-cuts. *Scand J For Res* 20:235–242
- Rudolphi J, Jonsson MT, Gustafsson L (2014) Biological legacies buffer local species extinction after logging. *J Appl Ecol* 51:53–62
- Runnel K, Rosenvald R, Löhms A (2013) The dying legacy of green-tree retention: Different habitat values for polypores and wood-inhabiting lichens. *Biol Conserv* 159:187–196
- Sahlén E, Ranius T (2009) Habitat availability in forests and clearcuts for saproxylic beetles associated with aspen. *Biodivers Conserv* 18:621–638
- Sahlén E, Schroeder LM (2010) Importance of habitat patch size for occupancy and density of aspen-associated saproxylic beetles. *Biodivers Conserv* 19:1325–1339
- Santaniello F, Djupström LB, Ranius T, Weslien J, Rudolphi J, Sonesson J (2017b) Simulated long-term effects of varying tree retention on wood production, dead wood and carbon stock changes. *J Environ Manage* 201:37–44
- Santaniello F, Djupström LB, Ranius T, Weslien J, Rudolphi J, Thor G (2017a) Large proportion of wood dependent lichens in boreal pine forest are confined to old hard wood. *Biodivers Conserv* 26:1295–1310
- Santaniello F, Line DB, Ranius T, Rudolphi J, Widenfalk O, Weslien J (2016) Effects of partial cutting on logging productivity, economic returns and dead wood in boreal pine forest. *For Ecol Manag* 365:152–158
- Schei FH, Blom HH, Gjerde I, Grytnes JA, Heegaard E, Sætersdal M (2013) Conservation of epiphytes: Single large or several small host trees? *Biol Conserv* 168:144–151
- Schroeder LM, Ranius T, Ekbohm B, Larsson S (2006) Recruitment of saproxylic beetles in high stumps created for maintaining biodiversity in a boreal forest landscape. *Can J For Res* 36:2168–2178
- Schroeder LM, Weslien J, Lindelöw Å, Lindhe A (1999) Attacks by bark- and wood-boring Coleoptera on mechanically created high stumps of Norway spruce in the two years following cutting. *For Ecol Manag* 123:21–30
- Shorohova E, Sinkevich S, Kryshen A, Vanha-Majamaa I (2019) Variable retention forestry in European boreal forests in Russia. *Ecol Process* 8:34
- Sira-Pietikainen A, Haimi J (2009) Changes in soil fauna 10 years after forest harvestings: Comparison between clear felling and green-tree retention methods. *For Ecol Manag* 258:332–338
- Simonsson P, Gustafsson L, Östlund L (2015) Retention forestry in Sweden: driving forces, debate and implementation 1968–2003. *Scand J For Res* 30:154–173
- Skogsstyrelsen (2019) Statistik om formellt skyddad skogsmark, frivilliga avsättningar, hänsynsytor samt improduktiv skogsmark Redovisning av regeringsuppdrag. Rapport 2019/18 – DNR 2018/4167. (in Swedish)
- Suominen M, Junninen K, Heikkala O, Kouki J (2015) Combined effects of retention forestry and prescribed burning on polypore fungi. *J Appl Ecol* 52:1001–1008
- Suominen M, Junninen K, Kouki J (2019) Diversity of fungi in harvested forests 10 years after logging and burning: Polypore assemblages on different woody substrates. *For Ecol Manag* 446:63–70
- Sverdrup-Thygeson A, Birkemoe T (2009) What window traps can tell us: effect of placement, forest openness and beetle reproduction in retention trees. *J Insect Conserv* 13:183–191
- Sverdrup-Thygeson A, Ims RA (2002) The effect of forest clearcutting in Norway on the community of saproxylic beetles on aspen. *Biol Conserv* 106:347–357
- Sveriges Officiella Statistik (2019) Statistiska Meddelanden. JO1403 SM 1901. Miljöhänsyn vid förnyingsavverkning Environmental consideration in regeneration felling. (in Swedish with English summary)
- Toivanen T, Heikkilä T, Koivula MJ (2014) Emulating natural disturbances in boreal Norway spruce forests: Effects on ground beetles (Coleoptera, Carabidae). *For Ecol Manag* 314:64–74
- Toivanen T, Kotiaho J (2007) Burning of logged sites to protect beetles in managed boreal forests. *Conserv Biol* 21:1562–1572
- Toivanen T, Kotiaho JS (2010) The preferences of saproxylic beetle species for different dead wood types created in forest restoration treatments. *Can J For Res* 40:445–464
- Vanha-Majamaa I, Lilja S, Ryömä R, Kotiaho JS, Laaka-Lindberg S, Lindberg H, Puttonen P, Tamminen P, Toivanen T, Kuuluvainen T (2007) Rehabilitating boreal forest structure and species composition in Finland through logging, dead wood creation and fire: The EVO experiment. *For Ecol Manag* 250:77–88
- Vanha-Majamaa I, Shorohova E, Kushnevskaya H, Jalonen J (2017) Resilience of understory vegetation after variable retention felling in boreal Norway spruce forests - A ten-year perspective. *For Ecol Manag* 393:12–28
- Weslien J, Djupström LB, Schroeder M, Widenfalk O (2011) Long-term priority effects among insects and fungi colonizing decaying wood. *J Anim Ecol* 80:1155–1162
- Westerfelt P, Weslien J, Widenfalk O (2018) Population patterns in relation to food and nesting resource for two cavity-nesting bee species in young boreal forest stands. *For Ecol Manag* 430:629–638
- Westerfelt P, Widenfalk O, Lindelöw Å, Gustafsson L, Weslien J (2015) Nesting of solitary wasps and bees in natural and artificial holes in dead wood in young boreal forest stands. *Insect Conserv Divers* 8:493–504
- Widerberg Koch M, Ranius T, Drobyshev I, Lindblad M (2012) Increased openness around retained oaks increases species richness of saproxylic beetles. *Biodivers Conserv* 21:3035–3059
- Wikberg S, Perhans K, Kindstrand C, Boberg Djupström L, Boman M, Mattsson L, Schroeder LM, Gustafsson L (2009) Cost-effectiveness of conservation strategies implemented in boreal forests: The area selection process. *Biol Conserv* 142:614–624

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.